BULL. AUSTRAL. MATH. SOC. 26A45
VOL. 11 (1974), 225-229.

An integral representation for a generalised variation of a function

A.M. Russell

In this note we present sufficient conditions for the continuity of the total kth variation of a function defined on a closed interval $[a, b]$. We also give an integral representation for total kth variation, thus obtaining an extension of the classical result

$$V(f;\ a,\ x) = \int_a^x |f'(t)|dt\ ,\quad a \le x \le b\ .$$

The results presented in this note will be a continuation of results obtained in [2], and unless otherwise stated, all definitions and notation will be taken from [2].

THEOREM 1. *If* $f \in BV_k[a, b]$, $k \ge 2$, *and* f *has a* $(k-1)$*th Riemann* derivative throughout* $[a, b]$ *(one-sided, of course, at* a *and* b *), then* $V_k(f;\ a,\ x)$ *is a continuous function of* x.

Proof. Let $\varepsilon > 0$ be arbitrary, and let $V_k(x)$ denote $V_k(f;\ a,\ x)$. Then there exists a π subdivision $a = y_0, y_1, \ldots, y_n = x$ of $[a, x]$ with $y_{n-1} = x'$ arbitrarily close to x, such that

$$\sum_{i=0}^{n-k} |Q_{k-1}(f;\ y_i,\ \ldots,\ y_{i+k-1}) - Q_{k-1}(f;\ y_{i+1},\ \ldots,\ y_{i+k})| > V_k(x) - \frac{\varepsilon}{2}\ ,$$

and

Received 17 May 1974.

$$|Q_{k-1}(f;\, y_{n-k}, \ldots, y_{n-1}) - Q_{k-1}(f;\, y_{n-k+1}, \ldots, y_n)| < \frac{\varepsilon}{2} .$$

Therefore,

$$\begin{aligned} V_k(x') &\geq \sum_{i=0}^{n-k-1} |Q_{k-1}(f;\, y_i, \ldots, y_{i+k-1}) - Q_{k-1}(f;\, y_{i+1}, \ldots, y_{i+k})| \\ &= \sum_{i=0}^{n-k} |Q_{k-1}(f;\, y_i, \ldots, y_{i+k-1}) - Q_{k-1}(f;\, y_{i+1}, \ldots, y_{i+k})| \\ &\qquad - |Q_{k-1}(f;\, y_{n-k}, \ldots, y_{n-1}) - Q_{k-1}(f;\, y_{n-k+1}, \ldots, y_n)| \\ &> V_k(x) - \frac{\varepsilon}{2} - \frac{\varepsilon}{2} = V_k(x) - \varepsilon . \end{aligned}$$

Consequently we have $V_k(x) \geq V_k(x') > V_k(x) - \varepsilon$, and so it follows that $V_k(x') \to V_k(x)$ as $x' \to x - 0$. Similarly it can be shown that $V_k(x') \to V_k(x)$ as $x' \to x + 0$, and this now completes the proof.

We make two observations.

The result of Theorem 1 is an extension of the classical case in which the total variation $V(f;\, a, x)$ of a continuous function f is continuous.

The hypothesis of existence of the $(k-1)$th Riemann* derivative is necessary, as shown by the following example. Let $f(x) = |x|$, $-1 \leq x \leq +1$. If $k = 2$, then $V_2(f;\, -1, 0) = 0$, whereas $V_2(f;\, -1, x) = 2$ for all $x > 0$.

We now obtain an integral representation for the total kth variation $V_k(f;\, a, x)$. This will be a generalisation of the classical result

$$V(f;\, a, x) = \int_a^x |f'(t)|\,dt ,$$

when f has an integrable derivative.

We make use of the following

THEOREM 2. *If f is a real-valued function whose kth derivative exists and is bounded on $[a, b]$, then $f \in BV_k[a, b]$.*

The proof follows readily from [*1*, Section 1.2].

THEOREM 3. *Let* f *be a function whose* k*th derivative is continuous in* $[a, b]$. *Then* $f \in BV_k[a, b]$, *and*

$$(k-1)!V_k(f;\ a,\ x) = \int_a^x |f^{(k)}(t)|dt\ ,\quad a \le x \le b\ .$$

Proof. It follows immediately from Theorem 2 that $f \in BV_k[a, b]$. Let us denote $V_k(f;\ a,\ x)$ by $V_k(a,\ x)$. Then, using [2, Theorem 7], and noting that f has a $(k-1)$th Riemann* derivative in $[a, b]$, we obtain

$$\frac{1}{h}\left[V_k(a,\ x+h)-V_k(a,\ x)\right] = \frac{1}{h}\,V_k(x,\ x+h) \quad \text{when} \quad h > 0\ .$$

Let $y_0, y_1, \ldots, y_n$ be a π subdivision of $[x, x+h]$ such that $x = y_0 < y_1 < \ldots < y_n = x + h$ and all sub-intervals $[y_i, y_{i+1}]$ are of equal length l , so that $nl = h$. Then, using [1, Section 1.2], we obtain

$$\begin{aligned}
V_k(x,\ x+h) &= \sup_{\pi} \sum_{i=0}^{n-k} (y_{i+k}-y_i)\,|Q_k(f;\ y_i,\ \ldots,\ y_{i+k})| \\
&\ge \frac{1}{k!} \sum_{i=0}^{n-k} (y_{i+k}-y_i)\,|f^{(k)}(\xi_i)|\ , \\
&\qquad\qquad \text{where } y_i < \xi_i < y_{i+k}\ ,\quad i = 0, 1, \ldots, k\ , \\
&\ge \inf_{x\le t\le x+h} |f^{(k)}(t)|\ \frac{1}{k!} \sum_{i=0}^{n-k} (kl) \\
&= \frac{h}{(k-1)!} \inf_{x\le t\le x+h} |f^{(k)}(t)|\cdot\left(1 - \frac{k-1}{n}\right)\ .
\end{aligned}$$

Letting n tend to infinity gives us the result

$$(1) \qquad V_k(x,\ x+h) \ge \frac{h}{(k-1)!} \inf_{x\le t\le x+h} |f^{(k)}(t)| \quad \text{when} \quad h > 0\ .$$

We now consider any $\pi(x_0, \ldots, x_n)$ subdivision of $[x, x+h]$ and use [1, Section 1.2] again to show that

$$\sum_{i=0}^{n-k} (x_{i+k}-x_i)\,|Q_k(f;\, x_i,\, \ldots,\, x_{i+k})| \le \sup_{x\le t\le x+h} |f^{(k)}(t)|\, \frac{1}{k!} \sum_{i=0}^{n-k} (x_{i+k}-x_i)$$
$$\le \frac{h}{(k-1)!} \sup_{x\le t\le x+h} |f^{(k)}(t)| \ .$$

Consequently,

$$V_k(x,\, x+h) \le \frac{h}{(k-1)!} \sup_{x\le t\le x+h} |f^{(k)}(t)| \ . \tag{2}$$

Combining inequalities (1) and (2) gives

$$\inf_{x\le t\le x+h} |f^{(k)}(t)| \le (k-1)!\, \frac{V_k(x,x+h)}{h} \le \sup_{x\le t\le x+h} |f^{(k)}(t)| \ . \tag{3}$$

Similar inequalities would be obtained when $h < 0$, so we conclude that $\frac{d}{dx} V_k(a,\, x)$ exists, and equals $\frac{1}{(k-1)!} |f^{(k)}(x)|$. The required result now follows.

REMARK. Continuity of $f^{(k)}$ in the preceding theorem is not necessary. If $f^{(k)}$ is bounded and continuous almost everywhere, then $V_k(f;\, a,\, x)$ is absolutely continuous by (3), and at a point x of continuity of $f^{(k)}$ we can show, as before, that

$$(k-1)!\, \frac{d}{dx} V_k(f;\, a,\, x) = |f^{(k)}(x)| \ .$$

Consequently, the integral representation still holds, the integral of course being Lebesgue.

We conclude with the following

THEOREM 4. *Let f be a function whose kth derivative is continuous. Then $f^{(k-s)} \in BV_s[a,\, b]$, $s = 1,\, 2,\, \ldots,\, k$, and*

$$(k-1)!\, V_k(f;\, a,\, x) = (s-1)!\, V_s\big(f^{(k-s)};\, a,\, x\big) \ , \quad a \le x \le b \ ,$$
$$s = 1,\, 2,\, \ldots,\, k \ .$$

Proof. It follows immediately from Theorem 3 that $f \in BV_k[a,\, b]$. That $f^{(k-s)} \in BV_s[a,\, b]$, $s = 1,\, 2,\, \ldots,\, k$ follows from [2, Theorem 12].

The required result now follows readily by an application of Theorem 3.

References

[1] L.M. Milne-Thomson, *The calculus of finite differences* (Macmillan & Co., London, 1933).

[2] A.M. Russell, "Functions of bounded kth variation", *Proc. London Math. Soc.* (3) **26** (1973), 547-563.

Department of Mathematics,
University of Melbourne,
Parkville,
Victoria.